

PURELL® Waterless Surgical Scrub Two-Application Method

DIRECTIONS FOR USE

Effective use of this product requires 2 mL per hand/arm (minimum two pump activations)

IF HANDS ARE VISIBLY SOILED, WASH HANDS FIRST WITH SOAP AND WATER.



STEP 1

- Remove debris from underneath fingernails using a disposable nail cleaner under running water¹
- Dry hands completely with a clean, disposable paper towel



APPLICATION 1 (Right Hand)

STEP 2

- Dispense 2 mL of product (dispenser will automatically actuate twice) into the palm of your right hand



STEP 3

- Dip fingertips of left hand into the product and work under nails



STEP 4

- Use the right hand to spread remaining product evenly over your left hand and forearm, to the elbow
- Pay attention to all skin surfaces including the nails, cuticles and web area in between fingers



APPLICATION 2 (Left Hand)

STEP 5

- Dispense 2 mL of product into palm of left hand
- Dip fingertips of right hand into the product and work under nails
- Use left hand and spread remaining product evenly over your right hand and forearm, to the elbow
- Pay attention to all skin surfaces including the nails, cuticles and web area in between fingers



STEP 6

- Allow hands and arms to completely air dry before donning surgical gown and gloves

IF DESIRED OR REQUIRED BY YOUR OPERATING ROOM PROTOCOL,
a third application can be used and spread evenly over both hands and forearms.



¹Healthcare personnel should keep natural fingernails no more than 2mm (0.08 inch) long according to AORN standards.